

# Acetaminophen (Tylenol®) Dosage Chart for Infants and Children

Dosing Instructions: Give every 4–6 hours as needed for fever or pain. Do NOT give more than 4 doses in 24 hours. Do NOT use with any other medicine containing acetaminophen. If possible, use weight to dose; otherwise, use age.

Age	0–3 months	4–11 months	12–23 months	2–3 years	4–5 years	6–8 years	9–10 years	11 years	12+ years
Weight (pounds)	6–11 lbs	12–17 lbs	18–23 lbs	24–35 lbs	36–47 lbs	48–59 lbs	60–71 lbs	72–95 lbs	96+ lbs
Infants Acetaminophen Liquid (160 mg / 5 mL)	1.25 mL	2.5 mL	3.75 mL	—	—	—	—	—	—
Children's Acetaminophen Liquid (160 mg / 5 mL)	—	—	—	5 mL	7.5 mL	10 mL	12.5 mL	15 mL	20 mL
Children's Acetaminophen Chewables (160 mg)	—	—	—	1 tablet	1½ tablets	2 tablets	2½ tablets	3 tablets	4 tablets
Adult's Acetaminophen Tablets (325 mg)	—	—	—	—	—	1 tablet	1 tablet	1½ tablets	2 tablets
Adult's Acetaminophen Tablets (500 mg)	—	—	—	—	—	—	—	1 tablet	1 tablet

Age Limits: Do NOT use acetaminophen for infants less than 12 weeks old unless your child's doctor tells you to.

Liquid Medicines: Use the syringe or dosing cup that comes with the medication to give an accurate dose.

Kitchen teaspoons or tablespoons should not be used.

See reverse side for ibuprofen dosing table.



# Ibuprofen (Motrin<sup>®</sup>, Advil<sup>®</sup>) Dosage Chart for Infants and Children

Dosing Instructions: Give every 6 hours as needed for fever or pain. Do NOT give more than 4 doses in 24 hours. Do NOT use with any other medicine containing ibuprofen. If possible, use weight to dose; otherwise, use age.

Age	6–11 months	12–23 months	2–3 years	4–5 years	6–8 years	9–10 years	11 years	12+ years
Weight (pounds)	12–17 lbs	18–23 lbs	24–35 lbs	36–47 lbs	48–59 lbs	60–71 lbs	72–95 lbs	96+ lbs
Infant's Ibuprofen Drops (50 mg / 1.25 mL)	1.25 mL	1.875 mL	2.5 mL	3.75 mL	5 mL	—	—	—
Children's Liquid Ibuprofen (100 mg / 5 mL)	2.5 mL	4 mL	5 mL	7.5 mL	10 mL	12.5 mL	15 mL	20 mL
Children's Ibuprofen Chewable Tablets (50 mg)	—	—	2 tablets	3 tablets	4 tablets	5 tablets	6 tablets	8 tablets
Junior Strength Ibuprofen Tablets (100 mg)	—	—	—	—	2 tablets	2 ½ tablets	3 tablets	4 tablets
Adult's Ibuprofen Tablets (200 mg)	—	—	—	—	1 tablet	1 tablet	1 ½ tablets	2 tablets

Age Limits: Do NOT use ibuprofen for infants less than 6 months old unless your child's doctor tells you to.

Liquid Medicines: Use the syringe or dosing cup that comes with the medication to give an accurate dose. Kitchen teaspoons or tablespoons should not be used.

See reverse side for acetaminophen dosing table.

BRV774808

